


Press Information

April 9, 2008

No. 3/2008 (46)



Max Planck Institute
for Chemical Ecology

At home on a crab, with new evolutionary neighbors

Scientists at the Max Planck Institute for Chemical Ecology, Germany, have rediscovered *Drosophila endobranchia*, a fly living in the mouth of land crabs

The members of *Drosophilidae*, a family consisting of about 3000 species, are often referred to as fruit flies although most of the members feed on microbes. As microbes can be found growing on a wide range of substrates, fruit flies can accordingly also be found in a multitude of habitats. One of the more bizarre choices of breeding substrates comes from *Drosophila endobranchia*. This species is one out of three known fruit flies that have found a home on (and inside) land-crabs. Although frequently mentioned in textbooks, the crab flies have surprisingly been neglected in active research since their description. *D. endobranchia* has actually not even been seen since its initial discovery in 1966. In January 2007, scientists from Bill Hansson's group at the Max Planck Institute for Chemical Ecology, Germany, managed to relocate these elusive flies on Grand Cayman in the Caribbean (the sole known home of this species), where small fragmented populations still persist in the few remaining patches of suitable habitats. Concomitant with the insects' re-exploration, a long-standing question regarding *D. endobranchia*'s evolutionary position within the *Drosophilidae*, disputed since its discovery due to a conflicting set of morphological characteristics, was resolved, based on a new comprehensive molecular and morphological analysis. (PLoS ONE, April 9, 2008)

Managing Director

Prof. Jonathan Gershenzon
Tel.: +49 (0)3641 - 57 1301
gershenzon@ice.mpg.de

Research Coordination

Dr. Jan-W. Kellmann
Tel.: +49 (0)3641 - 57 1000
Mobile: 0160 - 1622377
jkellmann@ice.mpg.de

Public Relations

Angela Overmeyer M.A.
Tel.: +49 (0)3641 - 57 2110
FAX: +49 (0)3641 - 57 1002
overmeyer@ice.mpg.de

Address

Beutenberg Campus
Hans-Knoell-Str. 8
07745 Jena

Internet

www.ice.mpg.de



MAX PLANCK SOCIETY



Courting of *Drosophila endobranchia* flies on their host crab. Picture: Max Planck Institute for Chemical Ecology/Stensmyr

Related pictures on request!

A well known species of the *Drosophilidae* family is the fruit fly, *Drosophila melanogaster*, model organism for genetic studies since many decades. There are also three members of the family, *D. endobanchia*, *D. carcinophila*, and *Lissocephala powelli*, which are known to fulfill all or parts of their life cycle on land crabs, and each of them in different areas of the globe. While *Lissocephala powelli* seems to be restricted to the Christmas Island in the Indian Ocean, the other two species appear at different locations of the Caribbean [1]. *D. endobanchia*, the least described of the three, was last seen on the Cayman Islands in 1966. The phylogenetic position of this species, i.e. the place it occupies in the tree of life, however, has remained disputed. Therefore, Marcus Stensmyr and colleagues decided to relocate this species in the Cayman Islands in order to re-define the phylogeny of this organism with current molecular techniques and shed light on the evolutionary history of this interesting trait.

A search for the Cayman crab flies in 2007 was successful, leading to the collection of 66 specimens. The phylogeny of this species was determined with the aid of molecular biology, which allowed the Max Planck researchers to place *D. endobanchia* within the *canalineae* species group, a little known Neotropical group of forest dwelling flies that belongs to the large *repleta* radiation that also includes *D. carcinophila*, the second Caribbean crab fly. Hence, both Caribbean crab fly species turned out to be quite closely related to each other, suggesting that something in their shared ancestry has made these flies more suitable for their most unconventional lifestyle, i.e. living on landcrabs. “Now it makes us wonder which aspects and traits of their shared ancestry allow them to survive and thrive in this most specialized environment”, says Marcus Stensmyr.

The Max Planck Institute for Chemical Ecology in Jena, Germany, consists of five departments constituting five independent working areas. The Department of Evolutionary Neuroethology by Prof. Bill S. Hansson was inaugurated in April 2006. It concentrates on the functional and evolutionary analysis of arthropod chemosensory systems. In addition, two departments focus on plant biology: Molecular Ecology (headed by Prof. Ian T. Baldwin) and Biochemistry (Prof. Jonathan Gershenzon). The department of Bioorganic Chemistry (Prof. Wilhelm Boland) specializes in chemosynthesis protocols and analytical techniques, and the department of Entomology (Prof. David G. Heckel) focuses on insect genomics. Two further Research Groups (Mass Spectrometry and Nuclear Magnetic Resonance Spectroscopy) support the scientists from all five departments with further analytical skills and measurement services.

[VG, JWK]

Citation: The Cayman Crab Fly Revisited—Phylogeny and Biology of *Drosophila endobanchia*. Stensmyr MC, Stieber R, Hansson BS. PLoS ONE 3(4), April 9, 2008. <http://www.plosone.org/doi/pone.0001942>

[1] Further reading: Flies' lives on a crab. Marcus C. Stensmyr, Bill S. Hansson. Current Biology 17, 743-746 (2007).

Contact: Bill S. Hansson, MPI for Chemical Ecology, Hans Knoell Str 8, 07745 Jena, Germany, +49 3641 57-1401, hansson@ice.mpg.de

Pictures: Angela Overmeyer M.A., MPI for Chemical Ecology, Hans Knoell Str 8, 07745 Jena, Tel.: +49 3641 57-2110, overmeyer@ice.mpg.de